

Supporting Online Information for ‘Archaeological Data Provide Alternative Hypotheses’ (McKechnie et al.)

Table S1. Archaeological data for Pacific herring and associated chronological information for all sites (N = 171). Relative abundance and rank order data based on number of identified specimens (NISP).

| No. | Region ^a | Sub-Region ^b | Arch. Site No. | Zooarchaeological Reference | NISP ^c | N Taxa | % NISP Herring | Rank Order ^d | Time Interval Years BP ^e | Reference for ¹⁴ C Date & stratigraphic context |
|-----|---------------------|-------------------------|----------------|--|-------------------|--------|----------------|-------------------------|-------------------------------------|--|
| 1 | SS | SVI | DcRu-75 | Wilson et al. 2003:85, Table 32 (1) | 682 | 13 | 42% | 2 | 300-1200 | Wilson et al. 2003:93 (1) |
| 2 | SS | SVI | DcRu-78 | Wigen 2009:3, Tables 2 & 3 (2) | 61 | 4 | 90% | 1 | Historic/300-1200 | Mitchell 1981 (3) |
| 3 | SS | SVI | DcRu-4 | Stewart & Wigen 2003:29, Table 1 (4) | 903 | 13 | 64% | 1 | 300-1200 | Gajewski et al. 2011 (5) |
| 4 | SS | SVI | DcRt-16 | Wilson et al. 2004:114, Table 45 (6) | 746 | 13 | 79% | 1 | 300-1200 | Wilson et al. 2004:121 (6) |
| 5 | SS | SVI | DcRt-10 | Willerton 2009:150, Table 12.4 (7) | 3167 | 13 | 77% | 1 | 300-1200/1200-2500 | Willerton 2009:36 (7) |
| 6 | SS | SVI | DcRu-71 | Wigen 2000:11, Table 5 (8) | 78 | 6 | 81% | 1 | 1200-2500/2500-5000* | Eldridge 2000:33 (9) |
| 7 | SS | SVI | DcRw-57 | Bowie & Wigen 2010:42, Table 2 (10) | 205 | 10 | 0.5% | 7 | 300-1200* | Bowie & Wigen 2010:48 (10) |
| 8 | SS | SVI | DcRv-21 | O'Neil 2007:40, Table 3 (11) | 57 | 5 | 77% | 1 | 300-1200* | O'Neil 2007:28 (11) |
| 9 | SS | SVI | DdRu-81 | Bowie et al. 2010:164, Table 29 (12) | 1815 | 20 | 75% | 1 | 300-1200 | Bowie et al. 2010:184 (12) |
| 10 | SS | SVI | DdRu-4 | Wilson 2005:154, Table 67 (13) | 638 | 14 | 62% | 1 | 300-1200 | Cooper 2008 (14); Wilson 2005 (13) |
| 11 | SS | SVI | DeRu-1 | Hewer 1999:26, Table 4 (15) | 1589 | 17 | 86% | 1 | 1200-2500 | Hewer 1999:52 (15) |
| 12 | SS | SVI | DdRu-5 | Kanipe et al. 2007:Appendix E, Table 11 (16) | 316 | 7 | 89% | 1 | 300-1200 | Kanipe et al. 2007:283-284 (16) |
| 13 | SS | SVI | DdRu-18 | Weathers et al. 2007:31, Table 8 (17) | 57 | 10 | 21% | 2 | 300-1200* | Weathers et al. 2007:36 (17) |
| 14 | SS | GI-BC | DdRt-6 | O'Neil & Wilson 2005:47, Table 6 (18) | 597 | 12 | 93% | 1 | 300-1200/1200-2500/2500-5000* | O'Neil & Wilson 2005:52 (18) |
| 15 | SS | GI-BC | DeRu-44 | Wilson 1988:Appendix 1 (19) | 93 | 4 | 90% | 1 | 1200-2500 | Wilson 1988:44 (19) |
| 16 | SS | GI-BC | DfRu-3 | Brolly et al. 1993:71, Table 7 (20) | 1155 | 9 | 88% | 1 | 300-1200/1200-2500 | Brolly et al. 1993:91-92 (20) |
| 17 | SS | GI-BC | DeRt-1 | Hanson 1995:36 & 39 (21) | 473 | 12 | 75% | 1 | 300-1200 | Hanson 1995:32 (21) |
| 18 | SS | GI-BC | DeRt-109 | Wigen 2007:103, Table 12 (22) | 521 | 13 | 85% | 1 | 300-1200 | Sumpter & Fedje 2007:62 (23) |
| 19 | SS | GI-BC | DgRu-3 | Ewonus, 2011:77-78, Table 1 (24) | 12675 | 29 | 70% | 1 | 1200-2500 | Grier 2006:102 (25) |

| No. | Region ^a | Sub-Region ^b | Arch. Site No. | Zooarchaeological Reference | NISP ^c | N Taxa | % NISP Herring | Rank Order ^d | Time Interval Years BP ^e | Reference for ¹⁴ C Date & stratigraphic context |
|-----|---------------------|-------------------------|----------------|--|-------------------|--------|----------------|-------------------------|-------------------------------------|--|
| 20 | SS | GI-BC | DgRv-4 | McLay 1999: Appendix 5 (26) | 1681 | 9 | 96% | 1 | 300-1200* | McLay 1999:67 (26) |
| 21 | SS | GI-BC | DgRv-1 | McLay 1999: Appendix 5 (26) | 246 | 9 | 86% | 1 | 300-1200* | McLay 1999:67 (26) |
| 22 | SS | GI-BC | DgRv-2 | McLay 1999: Appendix 5 (26) | 404 | 6 | 93% | 1 | 300-1200* | McLay 1999:67 (26) |
| 23 | SS | GI-BC | DgRv-2 | Matson et al. 1999:61-71 (27) | 2039 | 14 | 74% | 1 | 300-1200 | Matson 2003:85 (28) |
| 24 | SS | GI-BC | DgRw-78 | McLay 1999: Appendix 5 (26) | 59 | 6 | 39% | 1 | 300-1200* | McLay 1999:67 (26) |
| 25 | SS | GI-BC | DgRw-73 | McLay 1999: Appendix 5 (26) | 646 | 7 | 89% | 1 | 300-1200* | McLay 1999:67 (26) |
| 26 | SS | GI-BC | DgRw-32 | McLay 1999: Appendix 5 (26) | 80 | 8 | 55% | 1 | 300-1200* | McLay 1999:67 (26) |
| 27 | SS | GI-BC | DgRw-4 | Oliver & Park 1997:Appendix 3 (29) | 552 | 8 | 86% | 1 | 300-1200/1200-2500 | Burley 1989:33 (30) |
| 28 | SS | GI-BC | DgRw-204 | van Gaalen 1991:Table 3 (31) | 631 | 8 | 72% | 1 | 1200-2500 | Curtin 1991:38 (32) |
| 29 | SS | GI-BC | DiSe-10 | Eldridge 1987:5-63-65, Table 5-13-15 (33) | 166 | 8 | 71% | 1 | 2500-5000* | Eldridge 1987:5-14 (33) |
| 30 | SS | EVI | DjSf-13 | Mason & Hoffmann 1998:AIV-9, Table AIV-8 (34) | 221 | 7 | 92% | 1 | 300-1200/1200-2500 | Mason & Hoffmann 1998:Appendix V & VI (34); Mitchell 1974 (35) |
| 31 | SS | EVI | DkSf-26 | Mitchell 1988:17, Table 6 (36) | 4577 | 17 | 78% | 1 | 1200-2500/2500-5000 | Mitchell 1988:16 (36) |
| 32 | SS | EVI | DkSf-40 | Wilson & Clark 2003:40, Table 10 (37) | 83 | 5 | 83% | 1 | 300-1200 | Wilson & Clark 2003:47-48 (37) |
| 33 | SS | EVI | DkSf-4 | Simonsen 1991:Appendix tables (pg 53 of 59) (38) | 2612 | 10 | 89% | 1 | 1200-2500 | Simonsen 1991:27 (38) |
| 34 | SS | EVI | DkSf-19 | Caldwell 2008:120, Table 5.39 (39) | 2721 | 8 | 96% | 1 | 1200-2500 | Caldwell 2008:175 (39) |
| 35 | SS | EVI | DkSf-29 | Hall et al. 2003:83-93 (40) | 909 | 10 | 88% | 1 | 300-1200/1200-2500 | Hall et al. 2003:33 (40) |
| 36 | SS | EVI | DiSc-26 | Wilson 2003:42 Table 13 (41) | 215 | 6 | 86% | 1 | 300-1200 | Wilson 2003:52-53 (41) |
| 37 | SS | EVI | DiSe-7 | Wilson et al. 2004:44, Table 5 (42) | 5638 | 14 | 94% | 1 | 300-1200/1200-2500/2500-5000 | Wilson et al. 2004:48-50 (42) |
| 38 | SS | EVI | DhSb-3 | Wilson et al. 2006:60, Table 18 (43) | 661 | 9 | 97% | 1 | 1200-2500 | Wilson et al. 2006:83-84 (43) |
| 39 | SS | EVI | DhSb-11 | Bowie & Wigen 2010:41, Table 3 (10) | 2464 | 10 | 97% | 1 | 300-1200 | Kristensen et al. 2009:87, (44) |
| 40 | SS | EVI | DhRx-16 | Wilson & Crockford 1994:105, Table 38 (45) | 2870 | 10 | 97% | 1 | 1200-2500 | Wilson & Crockford 1994:v (45) |

| No. | Region ^a | Sub-Region ^b | Arch. Site No. | Zooarchaeological Reference | NISP ^c | N Taxa | % NISP Herring | Rank Order ^d | Time Interval Years BP ^e | Reference for ¹⁴ C Date & stratigraphic context |
|-----|---------------------|-------------------------|----------------|---|-------------------|--------|----------------|-------------------------|-------------------------------------|--|
| 41 | SS | EVI | DhRx-66 | Stryd et al. 1993:70 (46) | 2578 | 9 | 97% | 1 | 300-1200 | Stryd et al. 1993:81 (46) |
| 42 | SS | EVI | DhRx-104 | Pawlowski 2008:16, Table 4 (48) | 195 | 3 | 89% | 1 | 300-1200* | Pawlowski 2008:17 (48) |
| 43 | SS | EVI | DgRw-46 | Storey 2010:44, Table 6 (49) | 65 | 3 | 97% | 1 | 300-1200 | Storey 2010:37 (49) |
| 44 | SS | EVI | DgRw-34 | Willows 2009:39, Table 3 (50); Willows et al. 2012:118-119, Tables 14&15 (51) | 1668 | 17 | 87% | 1 | 300-1200/1200-2500 | Willows et al. 2012:138 (51) |
| 45 | SS | EVI | DeRv-68 | Brunsdon & Cooper 2008:17, Tables 2 & 3 combined (52) | 456 | 5 | 64% | 1 | 300-1200* | Brunsdon & Cooper 2008:10 (52) |
| 46 | SS | ESG | EaSe-18 | This Study | 5715 | 10 | 85% | 1 | 300-1200 | Springer et al. 2014 (53). |
| 47 | SS | ESG | EaSe-11 | This Study | 409 | 7 | 74% | 1 | 300-1200 | Springer et al. 2014 (53). |
| 48 | SS | ESG | EaSe-76 | This Study | 705 | 9 | 37% | 2 | 1200-2500 | Springer et al. 2014 (53). |
| 49 | SS | ESG | DISd-3 | This Study | 595 | 10 | 93% | 1 | 300-1200 | Springer et al. 2014 (53). |
| 50 | SS | ESG | DISd-6 | <i>Teeshoshum</i> - This Study | 1410 | 8 | 93% | 1 | 300-1200 | Springer et al. 2014 (53). |
| 51 | SS | ESG | DISe-10 | This Study | 64 | 6 | 9% | 4 | 300-1200 | Springer et al. 2014 (53). |
| 52 | SS | ESG | DISd-17 | This Study | 680 | 9 | 93% | 1 | Historic/300-1200 | Springer et al. 2014 (53). |
| 53 | SS | ESG | DjRw-1 | This Study | 1791 | 10 | 97% | 1 | 300-1200 | Coupland et al. in prep. |
| 54 | SS | FD | DhRr-18 | Coupland 1991:88, Table 3 (54) | 5429 | 11 | 82% | 1 | 1200-2500 | Coupland 1991:89 (54) Coupland 1989:71 (55) |
| 55 | SS | FD | DhRt-6 | Brolly & Muir 1993:51, Table 11 (56) | 897 | 11 | 42% | 1 | 2500-5000 | Brolly & Muir 1993:31 (56) |
| 56 | SS | FD | DhRr-6 | Pierson 2011:77-78, Appendix 1 (57) | 1650 | 28 | 30% | 2 | 300-1200/1200-2500/2500-5000 | Pierson 2011:27 (57) |
| 57 | SS | FD | DhRr-18 | Trost 2005:111, Appendix C (58) | 4640 | 19 | 57% | 1 | 300-1200* | Lepofsky et al. 2007:196 (59) |
| 58 | SS | FD | DhRq-1 | Pierson 2011:83-84, Appendix 1 (57) | 2098 | 22 | 40% | 1 | 300-1200/1200-2500 | Pierson 2011:31 (57) |
| 59 | SS | FD | DgRr-6 | Casteel 1976:85, Table 5-2 (60) | 304 | 8 | 1% | 4 | 1200-2500/2500-5000/5000-10000 | Matson 1976 (61) |
| 60 | SS | FD | DgRs-14 | Wigen 2002:1 (62) | 221 | 9 | 32% | 1 | 1200-2500 | Hammon 1986:97 (63) |
| 61 | SS | FD | DgRs-2 | Kusmer 1994a:133, Table 5-12 (64); 1994b:197, Table 4 (65) | 1244 | 9 | 25% | 3 | 300-1200/1200-2500/2500-5000 | Kusmer 1994b (65); Stryd et al. 1994: volume 2:120 (66) |
| 62 | SS | FD | DgRs-1 | Brolly et al. 1996:162 Table 13 (67) | 4589 | 10 | 22% | 2 | 300-1200/1200-2500/2500-5000 | Brolly et al. 1996:156, 180, & 209 (67) |

| No. | Region ^a | Sub-Region ^b | Arch. Site No. | Zooarchaeological Reference | NISP ^c | N Taxa | % NISP Herring | Rank Order ^d | Time Interval Years BP ^e | Reference for ¹⁴ C Date & stratigraphic context |
|-----|---------------------|-------------------------|----------------|---|-------------------|--------|----------------|-------------------------|-------------------------------------|--|
| 63 | SS | FD | DgRr-1 | Matson 1992 (68) in Butler and Campbell 2004:349 (69) | 25659 | 12 | 6% | 3 | 1200-2500/2500-5000 | Matson 2010:1-2 (70) |
| 64 | SS | PSnd | 45-KI-23 | Butler 1987:Table 10-1, pg. 10-2 (71) | 3999 | 16 | 0.1% | 13 | 300-1200 | Blukis Onat 1987:5-7 (72) |
| 65 | SS | PSnd | 45-KP-115 | Butler & Baker 2002:11, Table 2 (73) cited in Butler & Campbell 2004:349 (69) | 806 | 8 | 43% | 1 | 300-1200 | Lewarch et al. 2002:48 (74) |
| 66 | SS | PSnd | 45-SK-43 | Nelson et al. 2010:53, Table 7.5 (75) | 14800 | 11 | 25% | 2 | 300-1200/1200-2500/2500-5000 | Nelson et al. 2010a:33 (75) |
| 67 | SS | PSnd | 45-KI-437 | Kopperl & Butler 2002:107, Table 10.1 (76) | 5321 | 16 | 80% | 1 | Historic/300-1200 | Stein & Phillips 2002:58 (77) see also Kopperl 2001:3 (78) |
| 68 | SS | GI-WA | 45-SJ-24 | Pegg 1999:67, Table 3.1 (79) | 18654 | 11 | 68% | 1 | 300-1200 | Pegg 1999:19 (79) see also Stein et al. 2003:303 (80) |
| 69 | SS | GI-WA | 45-SJ-169 | Wigen 2003:283, Table D.7 (81) | 3223 | 24 | 52% | 1 | 300-1200/1200-2500/2500-5000 | Walker 2003:76, Table 6.4 (82) |
| 70 | SS | GI-WA | 45-SJ-280 | Kopperl 2007:16, Table 1 (83) | 2450 | 14 | 6% | 3 | 300-1200/1200-2500/2500-5000 | Bovy et al. 2007:20 (84) |
| 71 | SS | PSnd | 45IS2 | Trost et al. 2010:B8, Table B.10 (85) | 16154 | 31 | 3.80% | 6 | 300-1200 | Schalk & Nelson 2010:102, (86) |
| 72 | SS | PSnd | 45-IS-263 | Smith et al. 2011:23, Table 13 (87) | 401 | 15 | 6% | 5 | 300-1200 | Smith et al. 2011:l (87) |
| 73 | SS | PSnd | 45-IS-119 | Nelson et al. 2010:30, Table 6.2 (88) | 160 | 10 | 1% | 7 | 300-1200 | Nelson et al. 2010b:41 (88) |
| 74 | SS | GI-WA | 45-SJ-200 | Kopperl 2000:27 Table 1 in Kenady 2000 (89) | 126 | 11 | 25% | 1 | 300-1200* | NA |
| 75 | SS | GI-WA | 45-JS-252 | Kenady 2000:11, Table 9 (90) | 373 | 6 | 19% | 2 | 300-1200 | Kenady 2000:Appdx. A (90) |
| 76 | SS | PSnd | 45-KI-428 & 29 | Wigen 1995:A5-66, A5-70, Table A5-13 (91) | 8057 | 29 | 2% | 8 | 300-1200/1200-2500/2500-5000 | Larson & Lewarch 1995: Table 6.1, pg 6-4-6-5 (92) |
| 77 | SS | PSnd | 45-PI-974 | Shantry et al. 2010:54, Table 13 (93) | 1226 | 12 | 43% | 1 | 300-1200 | Shantry et al. 2010 (93) |
| 78 | SS | PSnd | 45-MS-50 | Wessen 2004:46-47, Tables 14 & 15 (94) | 1248 | 7 | 2% | 7 | 300-1200/1200-2500 | Wessen 2004:Appendix B (94) |
| 79 | WCVI | WCVI | EdSv-4 | Wigen 1999:Table 3 (95) | 183 | 14 | 38.80% | 1 | Historic/300-1200/1200-2500* | NA |
| 80 | WCVI | WCVI | DiSo-9 | Calvert 1980:298 & 302, Tables 45 & 49 (96) | 19426 | 22 | 48% | 1 | 1200-2500 | Calvert 1980:123, Table 9 (96) |
| 81 | WCVI | WCVI | DgSI-67 | Wilson 1994:18, Table 2 (97) | 68 | 10 | 57% | 1 | 300-1200 | Wilson 1994:27 (97) |
| 82 | WCVI | BS | DfSj-40 | Brolly & Pegg 1998:167 (98) | 757 | 12 | 79% | 1 | 300-1200/1200-2500 | Brolly & Pegg 1998:156 (98) |

| No. | Region ^a | Sub-Region ^b | Arch. Site No. | Zooarchaeological Reference | NISP ^c | N Taxa | % NISP Herring | Rank Order ^d | Time Interval Years BP ^e | Reference for ¹⁴ C Date & stratigraphic context |
|-----|---------------------|-------------------------|----------------|---|-------------------|--------|----------------|-------------------------|-------------------------------------|---|
| 83 | WCVI | BS | DfSj-100 | Weathers et al. 2008:45, Table 6 (99); Wigen 2008 (100) | 1021 | 15 | 79% | 1 | 2500-5000 | Stryd et al. 1991:37 (101) |
| 84 | WCVI | BS | DfSj-57 | Spady & Wigen 2008:49, Table 8 (102) | 1852 | 14 | 84% | 1 | 300-1200 | Spady & Wigen 2008:60 (102) |
| 85 | WCVI | BS | DfSi-16 & 17 | McKechnie 2005a:125, Appendix B1 (103) | 6979 | 21 | 53% | 1 | 300-1200/1200-2500/2500-5000 | McMillan & St. Claire 2005:45&77 (104) |
| 86 | WCVI | BS | DfSi-26 | McKechnie 2005b:6, Table 2 (105) | 157 | 7 | 58% | 1 | 300-1200 | McKechnie, 2005b:5 (105) |
| 87 | WCVI | BS | 129T | This Study | 1738 | 17 | 61% | 1 | Historic/300-1200 | Sumpter & St. Claire 2009:112 (106); McKechnie 2013:255-256 (107) |
| 88 | WCVI | BS | 83T | This Study | 1997 | 16 | 47% | 1 | Historic/300-1200 | McKechnie 2013:255-256 (107) |
| 89 | WCVI | BS | 82T | This Study | 1374 | 16 | 39% | 2 | 300-1200 | McKechnie 2013:255-256 (107) |
| 90 | WCVI | BS | 306T | Wigen 2013:7, Table 3 (108) and This Study | 482 | 10 | 72% | 1 | Historic/300-1200 | Smith et al. 2012 (109) |
| 91 | WCVI | BS | 304T | This Study | 1155 | 14 | 41% | 2 | Historic/300-1200 | McKechnie 2013:255-256 (107) |
| 92 | WCVI | BS | 206T | This Study | 1939 | 17 | 41% | 1 | Historic/300-1200/1200-2500 | McKechnie 2013:255-256 (107) |
| 93 | WCVI | BS | 72T | Wigen 2009:Table 6 in Sumpter & St. Claire 2009:132 (2) | 3725 | 18 | 85% | 1 | 300-1200 | Sumpter and St. Claire 2009:84 (2) |
| 94 | WCVI | BS | DeSh-1 | Wigen 2003:Table 6 in Sumpter 2003:4.6 (110) | 700 | 17 | 48% | 1 | Historic/300-1200 | Sumpter 2003:18 (111) |
| 95 | WCVI | BS | DeSh-2 | Wigen 2003:Table 9 (110) | 169 | 6 | 18% | 2 | Historic/300-1200 | Sumpter 2003:18 (111) |
| 96 | WCVI | BS | DfSh-7 | McKechnie 2012: Table 2 (112) | 32401 | 33 | 81% | 1 | 300-1200/1200-2500/2500-5000 | McMillan & St. Claire 2012:33&93 (113) |
| 97 | WCVI | WCVI | DeSf-6 | McKechnie 2007a:7 (114) | 258 | 10 | 11% | 3 | 300-1200 | McKechnie & Fedje 2010 (115) |
| 98 | CC | NVI | EeSu-13 | Wilson et al. 1993:32, Table 6 (116) | 1121 | 15 | 67% | 1 | 2500-5000 | Wilson et al. 1993:69 (116) |
| 99 | CC | Namu | EISx-3 | Cannon et al. 2011:62, Table 5.1 (117) | 3356 | 14 | 48% | 2 | 300-1200/1200-2500 | Cannon 1997:15 (118) |
| 100 | CC | Namu | EISx-5 | Cannon et al. 2011:62, Table 5.1 (117) | 989 | 8 | 61% | 1 | 300-1200/1200-2500/2500-5000 | Cannon 2000:74 (119); Cannon 1998:40 (120) |
| 101 | CC | Namu | EISx-18 | Cannon et al. 2011:62, Table 5.1 (117) | 567 | 9 | 48% | 2 | 300-1200/1200-2500/2500-5000 | Cannon 1998:37 (120) |
| 102 | CC | Namu | EISx-10 | Cannon et al. 2011:62, Table 5.1 (117) | 1433 | 9 | 62% | 1 | 300-1200/1200-2500/2500-5000 | Cannon 1997:45 (118); Cannon 2000:72 (119) |

| No. | Region ^a | Sub-Region ^b | Arch. Site No. | Zoarchaeological Reference | NISP ^c | N Taxa | % NISP Herring | Rank Order ^d | Time Interval Years BP ^e | Reference for ¹⁴ C Date & stratigraphic context |
|-----|---------------------|-------------------------|--------------------|---|-------------------|--------|----------------|-------------------------|---|--|
| 103 | CC | Namu | EITa-25 | Cannon et al. 2011:62, Table 5.1 (117) | 105 | 6 | 35% | 2 | Historic/300-1200/1200-2500/2500-5000 | Cannon 1997:45 (118); Cannon 2000:72 (119) |
| 104 | CC | Namu | EISx-1 | Cannon et al. 2011:62, Table 5.1 (117) | 5870 | 15 | 41% | 2 | 300-1200/1200-2500/2500-5000/5000-10000 | Carlson 1991 (121) |
| 105 | CC | Namu | EkSx-12 (Koeye R.) | Cannon et al. 2011:62, Table 5.1 (117) | 1992 | 18 | 9.94% | 2 | 300-1200/1200-2500 | Cannon 1997:45 (118) |
| 106 | CC | Namu | EITa-3 | Cannon et al. 2011:62, Table 5.1 (117) | 137 | 5 | 12% | 3 | Historic/300-1200 | Cannon 2000:72 (119); Cannon 1997:45 (118) |
| 107 | CC | Namu | EITb-1 (Nulu) | Cannon et al. 2011:62, Table 5.1 (117) | 4802 | 13 | 86% | 1 | Historic/300-1200/1200-2500 | Cannon 2000:72 (119); Cannon 1998:22 (120) |
| 108 | CC | Namu | EITb-2 | Cannon et al. 2011:62, Table 5.1 (117) | 97 | 8 | 38% | 2 | 300-1200/1200-2500 | Cannon 1998:26 (120) |
| 109 | CC | Rint | EkSt-1 | Cannon et al. 2011:62, Table 5.1 (117) | 2535 | 5 | 29% | 2 | Historic/300-1200 | MacDonald et al. 2011:3623 (122) |
| 110 | CC | Rint | EjSw-1 | Cannon et al. 2011:62, Table 5.1 (117) | 3113 | 7 | 47% | 1 | 1200-2500 | MacDonald et al. 2011:3623 (122) |
| 111 | CC | Rint | EjSv-1 | Cannon et al. 2011:62, Table 5.1 (117) | 122 | 3 | 35% | 1 | 300-1200 | MacDonald 2008:26 (123) |
| 112 | CC | Rint | EjSv-2 | Cannon et al. 2011:62, Table 5.1 (117) | 374 | 6 | 56% | 1 | 300-1200/1200-2500/2500-5000 | MacDonald 2008:26 (123) |
| 113 | CC | Rint | EjSv-3 | Cannon et al. 2011:62, Table 5.1 (117) | 140 | 3 | 40% | 1 | 300-1200 | MacDonald et al. 2011:3623 (122) |
| 114 | CC | Rint | EjSv-4 | Cannon et al. 2011:62, Table 5.1 (117) | 64 | 5 | 56% | 1 | 1200-2500 | MacDonald 2008:26 (123) |
| 115 | CC | Rint | EjSv-5 | Cannon et al. 2011:62, Table 5.1 (117) | 328 | 4 | 66% | 1 | 1200-2500/2500-5000 | MacDonald et al. 2011:3623 (122) |
| 116 | CC | Rint | EjSv-8 | Cannon et al. 2011:62, Table 5.1 (117) | 222 | 4 | 48% | 1 | 300-1200/1200-2500 | MacDonald 2008:26 (123) |
| 117 | CC | Rint | EjSv-9 | Cannon et al. 2011:62, Table 5.1 (117) | 551 | 5 | 42% | 1 | 300-1200/1200-2500/2500-5000 | MacDonald 2008:26 (123) |
| 118 | CC | Rint | EjSv-10 | Cannon et al. 2011:62, Table 5.1 (117) | 905 | 7 | 52% | 1 | 300-1200/1200-2500/2500-5000 | MacDonald 2008:26 (123) |
| 119 | CC | Rint | EjSv-11 | Cannon et al. 2011:62, Table 5.1 (117) | 127 | 5 | 42% | 2 | 300-1200/1200-2500/2500-5000 | MacDonald 2008:26 (123) |
| 120 | HG | S. HG | FaTt-9 | Wigen 1990 (124); Acheson 1998 (125) | 23469 | 15 | 16% | 2 | 300-1200 | Acheson 1998 (125) |
| 121 | HG | S. HG | 1325T | Fedje et al. 2005:195, Table 11.2 (126) | 751 | 11 | 0.10% | 7 | 5000-10000 | Fedje et al. 2001 (127); Fedje et al. 2005 (126) |

| No. | Region ^a | Sub-Region ^b | Arch. Site No. | Zooarchaeological Reference | NISP ^c | N Taxa | % NISP Herring | Rank Order ^d | Time Interval Years BP ^e | Reference for ¹⁴ C Date & stratigraphic context |
|-----|---------------------|-------------------------|----------------|---|-------------------|--------|----------------|-------------------------|---------------------------------------|--|
| 122 | HG | S. HG | 2008T | Orchard 2011:23, Table 12 (128) | 254 | 9 | 1.60% | 5 | 300-1200* | Orchard 2011:22 (128) |
| 123 | HG | S. HG | 699T | Orchard 2007:236, Table 7.10 (129) | 11651 | 12 | 28% | 2 | Historic/300-1200 | Orchard 2007:223-227 (129) |
| 124 | HG | S. HG | 740T | Orchard 2007:236, Table 7.10 (129) | 112 | 10 | 13% | 3 | 300-1200 | Orchard 2007:223-227 (129) |
| 125 | HG | S. HG | 717T | Orchard 2007:236, Table 7.10 (129) | 6877 | 10 | 43% | 2 | 300-1200 | Orchard 2007:223-227 (129) |
| 126 | HG | S. HG | 924T | This study and Orchard 2011:9-10, Table 3&4 (128) | 35631 | 13 | 64.50% | 1 | Historic/300-1200/1200-2500/2500-5000 | Orchard 2007:258, Table 7.16 (129); Fedje et al. 2011:37 (130) |
| 127 | HG | S. HG | 1370T | Orchard 2011:17, Table 8 (128) | 149 | 8 | 24.20% | 2 | 300-1200 | Fedje & Smith 2009:5 (131) |
| 128 | HG | S. HG | 781T | Orchard 2007:236, Table 7.10; (129) combined with Orchard 2008:51-67, Tables D1-D10 & C1-C9 (132) | 3736 | 13 | 80% | 1 | Historic/300-1200 | Orchard 2008:27, Table 3 (132) |
| 129 | HG | S. HG | 922T | Sumpter 1999 (133); Wigen 1999 (134) | 1927 | 13 | 2% | 6 | 300-1200/1200-2500 | Sumpter 1999:1 (133) |
| 130 | HG | S. HG | 785T | Orchard 2007:236, Table 7.10 (129) | 6317 | 14 | 3% | 4 | Historic/300-1200/1200-2500 | Orchard 2007:223-227 (129) |
| 131 | HG | S. HG | 1134T | Orchard 2007:237, Table 7.10 (129) | 904 | 5 | 0.00% | Absent | Historic | Orchard 2007:223-227 (129) |
| 132 | HG | S. HG | 1127T | Steffen 2006:65, Table 3.6 (135) | 1615 | 11 | 0.43% | 8 | 5000-10000 | Fedje et al. 2005:209 (136) |
| 133 | HG | N. HG | FjUb-10 | Christensen & Stafford 2005 (137) | 1731 | 9 | 2% | 6 | 2500-5000/5000-10000 | Christensen & Stafford 2005:251 (137) |
| 134 | HG | N. HG | GaUa-18 | Christensen et al. 2010:69, Table 11 (138) | 1020 | 14 | 6% | 5 | 300-1200/1200-2500 | Christensen et al. 2010:54 (138) |
| 135 | NC | Dund | GdTq-1 | Brewster & Martindale 2011:258, Table 15.4 (139) | 379 | 4 | 4% | 3 | 2500-5000/5000-10000 | Brewster & Martindale 2011:256-7 (139) |
| 136 | NC | Dund | GcTr-5 | Brewster & Martindale 2011:258, Table 15.4 (139) | 671 | 4 | 7% | 3 | 1200-2500/2500-5000 | Brewster & Martindale 2011:256-7 (139) |
| 137 | NC | Dund | GcTq-5 | Brewster & Martindale 2011:258, Table 15.4 (139) | 896 | 4 | 11% | 2 | 1200-2500/2500-5000/5000-10000 | Brewster & Martindale 2011:256-7 (139) |
| 138 | NC | Dund | GcTq-7 | Brewster & Martindale 2011:258, Table 15.4 (139) | 204 | 4 | 12% | 3 | 300-1200/1200-2500/2500-5000* | Martindale et al. 2010:73 (140) |
| 139 | NC | Dund | GcTr-10 | Brewster & Martindale 2011:258, Table 15.4 (139) | 500 | 4 | 5% | 3 | 300-1200/1200-2500/2500-5000* | Martindale et al. 2010:73 (140) |
| 140 | NC | Dund | GcTr-8 | Brewster & Martindale 2011:258, Table 15.4 (139) | 347 | 4 | 26% | 2 | 1200-2500/2500-5000/5000-10000 | Brewster & Martindale 2011:256-7, Table 15.3 |
| 141 | NC | Dund | GcTr-9 | Brewster & Martindale 2011:258, Table 15.4 (139) | 75 | 4 | 19% | 2 | 300-1200/1200-2500/2500-5000* | Brewster & Martindale 2011:255 (139) |

| No. | Region ^a | Sub-Region ^b | Arch. Site No. | Zooarchaeological Reference | NISP ^c | N Taxa | % NISP Herring | Rank Order ^d | Time Interval Years BP ^e | Reference for ¹⁴ C Date & stratigraphic context |
|-----|---------------------|-------------------------|----------------|--|-------------------|--------|----------------|-------------------------|-------------------------------------|--|
| 142 | NC | Dund | GcTq-1 | Brewster & Martindale 2011:258, Table 15.4 (139) | 382 | 4 | 13% | 3 | 2500-5000/5000-10000 | Brewster & Martindale 2011:256-7 (139) |
| 143 | NC | Dund | GcTq-6 | Brewster & Martindale 2011:258, Table 15.4 (139) | 82 | 4 | 16% | 2 | 2500-5000/5000-10000 | Brewster & Martindale 2011:256-7 (139) |
| 144 | NC | Dund | GcTq-4 | Brewster & Martindale 2011:258, Table 15.4 (139) | 64 | 4 | 36% | 2 | 2500-5000/5000-10000 | Brewster & Martindale 2011:256-7 (139) |
| 145 | NC | PRH | GbTo-24 | This Study | 142 | 7 | 4% | 4 | 300-1200/1200-2500/2500-5000* | Martindale in prep. |
| 146 | NC | PRH | GcTo-6 | This Study | 155 | 5 | 10.30% | 2 | 1200-2500 | Coupland et al. 1993 (141) |
| 147 | NC | PRH | GbTo-77 | Coupland et al. 2010:199, Table 12 (142) | 83 | NA | 24.10% | 2 | 1200-2500/2500-5000 | Coupland et al. 2010:193 (142) |
| 148 | NC | PRH | GbTo-28 | Coupland et al. 2010:198, Table 8 (142) | 742 | NA | 21.20% | 2 | 1200-2500 | Coupland et al. 2010:193 (142) |
| 149 | NC | PRH | GbTo-46 | Coupland et al. 2010:198, Table 8 (142) | 419 | NA | 11.00% | 2 | 1200-2500 | Coupland et al. 2010:193 (142) |
| 150 | NC | PRH | GbTo-31 | Coupland et al. 2010:197, Table 6 (142) | 374 | NA | 3.20% | 2 | 300-1200/1200-2500 | Stewart et al. 2009 (143) |
| 151 | NC | PRH | GbTo-13 | Eldridge et al. 2008:117, Table 111 (144) | 284 | 7 | 16% | 2 | 1200-2500 | Eldridge et al. 2008:241 (144) |
| 152 | NC | PRH | GbTo-37 | Eldridge et al. 2008:117 (144) | 58 | 4 | 12% | 2 | 2500-5000 | Eldridge et al. 2008:241 (144) |
| 153 | SEAK | Angn | 49-SIT-124 | Moss 1989:242-260 (145) combined with Moss 2007:8-10, Unit 1 (146) | 1655 | 14 | 12% | 2 | 300-1200/1200-2500 | Moss 1989:249 (145) |
| 154 | SEAK | Angn | 49-SIT-130 | Moss 1989:299-309 (145) | 1042 | 2 | 0% | Ab-sent | 300-1200 | Moss 1989:299 (145) |
| 155 | SEAK | Angn | 49-SIT-132 | Moss 1989:203:Table 5.12 (145) | 356 | 6 | 41% | 2 | 300-1200 | Moss 1989:197 (145) |
| 156 | SEAK | Angn | 49-SIT-171 | Moss 1989: 214-224 (145) | 384 | 3 | 4% | 2 | 300-1200 | Moss 1989:217 (145) |
| 157 | SEAK | Angn | 49-SIT-244 | Moss 1989:175-177 (145) | 1730 | 9 | 55% | 1 | 300-1200 | Moss 1989:169 (145) |
| 158 | SEAK | Angn | 49-SIT-259 | Moss 1989, p. 267-269 (145) | 63 | 5 | 3% | 4 | 300-1200/1200-2500 | Moss 1989:227 (145) |
| 159 | SEAK | Angn | 49-SIT-299 | Moss 1989 p. 326-334 (145) | 1109 | 9 | 1% | 2 | 300-1200/1200-2500 | Moss 1989:324 (145) |
| 160 | SEAK | Angn | 49-SIT-304 | Moss 1989 p. 273-282 (145) | 920 | 2 | 99.70% | 1 | 300-1200 | Moss 1989:277 (145) |
| 161 | SEAK | Noyes | 49-CRG-188 | Moss 2011:160 (147) | 112 | 8 | 36.60% | 1 | 300-1200/1200-2500 | Moss 2004 (148) |

| No. | Region ^a | Sub-Region ^b | Arch. Site No. | Zooarchaeological Reference | NISP ^c | N Taxa | % NISP Herring | Rank Order ^d | Time Interval Years BP ^e | Reference for ¹⁴ C Date & stratigraphic context |
|-----|---------------------|-------------------------|----------------|--|-------------------|--------|----------------|-------------------------|-------------------------------------|--|
| 162 | SEAK | Suem | 49CRG-88 | Moss & Erlandson 2001:36-37, 40-41 (149) | 144 | 9 | 1% | 7 | 300-1200 | Moss & Erlandson 2001:35 (149) |
| 163 | SEAK | Dall | 49-DIX-46 | Moss 2008:50, Table 5 (150) | 194 | 12 | 5% | 4 | 1200-2500/2500-5000 | Moss 2008:49 (150) |
| 164 | SEAK | Annt | 49-KET-229 | Minor et al. 1986b (151) | 618 | 8 | 17% | 3 | 1200-2500 | Minor et al. 1986 (151) |
| 165 | SEAK | POW | 49-PET-67 | Moss 2011:160 (147) | 391 | 6 | 23% | 2 | 2500-5000 | Moss 2011b:160 (147) |
| 166 | SEAK | POW | 49-PET-556 | Moss 2011:160 (147) | 233 | 9 | 65 | 1 | 1200-2500/2500-5000 | Moss 2011b:160 (147) |
| 167 | SEAK | Kuiu | 49-XPA-29 | Maschner 1992:318-320 (152) | 10912 | 9 | 32% | 2 | 300-1200/1200-2500 | Maschner 1992:178 (152) |
| 168 | SEAK | Kuiu | 49-XPA-39 | Maschner 1992:318-320 (152) | 11726 | 7 | 66% | 1 | 300-1200/1200-2500/2500-5000 | Maschner 1992:178 (152) |
| 169 | SEAK | Kuiu | 49-XPA-106 | Maschner 1992:318-320 (152) | 515 | 4 | 3% | 3 | 300-1200 | Maschner 1992:178 (152) |
| 170 | SEAK | Kuiu | 49-XPA-112 | Maschner 1992:318-320 (152) | 548 | 4 | 1% | 2 | 300-1200 | Maschner 1992:178 (152) |
| 171 | SS | EVI | DhRx-101 | Eldridge et al. 2007:101-102, Table 8 (47) | 430 | 8 | 90% | 1 | 300-1200/1200-2500 | Eldridge et al. 2007:54 (47) |

a -Regions: SS = Salish Sea, Including Puget Sound, Strait of Georgia, & the Strait of Juan de Fuca; WCVI = Western Vancouver Island; CC = Central BC Coast; HG = Haida Gwaii; NC = Northern BC Coast; SEAK = Southeast Alaska.

b - Sub-Regions: PSnd = Puget Sound; GI-WA = Washington State Gulf Islands; GI-BC = BC Gulf Islands; SVI = Southern Vancouver Island; EVI = Eastern Vancouver Island; ESG = Eastern Strait of Georgia; FD = Fraser Delta; BS = Barkley Sound; WCVI = Western Vancouver Island; NVI = Northern Vancouver Island; Namu = Namu Area; Rinlt = Rivers Inlet Area; S. HG = Southern Haida Gwaii; N. HG = Northern Haida Gwaii; Dund = Dundas Archipelago; PRH = Prince Rupert Harbour; Angn = Angoon Area; Noyes = Noyes Island; Suemez = Suemez Island; Dall = Dall Island; Annt = Annette Island; POW = Prince of Wales Island; Kuiu = Kuiu Island

c – NISP = Number of identified specimens (fish only).

d – Rank Order = Rank order of herring based on numbers of identified specimens (NISP). A rank of ‘1’ indicates herring has the highest rank based on NISP. The higher the number the lower the rank.

e – Dates fall within these broad temporal intervals. Asterisk indicates no available ¹⁴C dates and where site chronology has been estimated by excavators based on the presence of temporally diagnostic artifacts (and the absence of modern ‘historic’ artifacts), and the relative Holocene sea level history for the region.