

CORRECTION

Open Access

Correction: A glycine zipper motif mediates the formation of toxic beta-amyloid oligomers in vitro and in vivo

Virginia Fonte¹, Vishantie Dostal¹, Christine M Roberts¹, Patrick Gonzales¹, Pascale N Lacor², Pauline T Velasco², Jordi Magrane³, Natalie Dingwell⁴, Emily Y Fan⁴, Michael A Silverman⁴, Gretchen H Stein⁵ and Christopher D Link^{1,6*}

Correction

After publication of this work [1], we noted that we inadvertently failed to include the complete list of all co-authors. The full list of authors has now been added and the Authors' contributions section has been modified accordingly.

Authors' contributions

C. elegans experiments were performed by VF, VD, CMR, PG, and CDL. Neuro 2a experiments were performed by GHS. Characterization of ADDL preps was performed by PTV. ADDL-binding and toxicity assays in hippocampal cultures were performed and analyzed by PNL, and Tau phosphorylation analysis done by ND, EYF and MAS. Adenovirus transfection of cortical neurons was performed by JM. GHS, PNL, MAS, JM and CDL prepared the manuscript. All authors have read and approved the final manuscript.

Acknowledgements

The authors would like to thank Hilary Devlin for assistance with media preparation. Some nematode strains were provided by the *Caenorhabditis* Genetics Center, funded by the NIH National Center for Research Resources. This work was supported by an Alzheimer's Association Zenith Award (C.D.L), NIH award AG12423 (C.D.L), and Canadian Institutes for Health Research, Grant # 102686 (M.A.S.). PNL and PTV were supported by funding granted to William Klein (Alzheimer's Association IIRG-06-26989; NIH award R01AG022547; AHA# A2006-092 and gift from Acumen Pharmaceuticals Inc).

Author details

¹Institute for Behavioral Genetics, University of Colorado, Boulder, CO 80309, USA. ²Department of Neurobiology and Physiology, Northwestern University, Evanston, IL 60208, USA. ³Department of Neurology and Neuroscience, Weill Medical College of Cornell University, New York, NY 10065, USA.

⁴Department of Biological Sciences, Simon Fraser University, Burnaby, BC V5A 1S6, Canada. ⁵Department of Molecular, Cellular and Developmental Biology, University of Colorado, Boulder, CO 80309, USA. ⁶Integrative Physiology, University of Colorado, Boulder, CO 80309, USA.

Received: 25 February 2014 Accepted: 25 February 2014

Published: 25 March 2014

* Correspondence: linkc@colorado.edu

¹Institute for Behavioral Genetics, University of Colorado, Boulder, CO 80309, USA

⁶Integrative Physiology, University of Colorado, Boulder, CO 80309, USA
Full list of author information is available at the end of the article

Reference

1. Fonte V, Dostal V, Roberts CM, Gonzales P, Lacor P, Magrane J, Dingwell N, Fan EY, Silverman MA, Stein GH, Link CD: A glycine zipper motif mediates the formation of toxic β -amyloid oligomers *in vitro* and *in vivo*. *Mol Neurodegener* 2011, 6:61.

doi:10.1186/1750-1326-9-12

Cite this article as: Fonte et al.: Correction: A glycine zipper motif mediates the formation of toxic beta-amyloid oligomers in vitro and in vivo. *Molecular Neurodegeneration* 2014 9:12.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

